

On twinning in smectic crystals

V.I. Marchenko

Kapitza Institute for Physical Problems, RAS, Moscow, 119334 Russia

It is shown that mechanical twinning in smectic crystals is possible. The structure of the boundary of twins for a small disorientation of crystallites is determined. The periodic twin structure, which should appear at the tension of the smectic layer, is proposed.

PACS numbers:

Keywords:

The energy of small deformations of a smectic crystal is given by the expression [1]

$$\mathcal{E} = \int \frac{A}{2} \left\{ \left(\partial_z u - \frac{(\partial_\alpha u)^2}{2} \right)^2 + \lambda^2 (\Delta_\perp u)^2 \right\} dV, \quad (1)$$

where u – is the displacement of layers along z axis (in the initial homogeneous undeformed state of the smectic crystal, layers lie in the xy plane, A – is the elastic modulus, λ – is the length parameter, ∂_α – is the gradient vector in the xy plane, and $\Delta_\perp = \partial_\alpha^2$).


According to Eq. (1), the undeformed state turned by small angle $\theta \ll 1$ in the xz plane (in this case, $\partial_x u = \theta$) corresponds to the derivative $\partial_z u = \theta^2/2$. Let us consider the boundary between the states $\partial_x u = \pm\theta$ ($x \rightarrow \pm\infty$) that lies in the yz plane. The quantity $\partial_z u$ is unchanged inside the boundary. The variation equilibrium equation in the problem under consideration reduces to the form

$$\lambda^2 f''' + \frac{\theta^2}{2} f' - \frac{3}{2} f^2 f' = 0, \quad (2)$$

where $f = \partial_x u$. The solution of this equation has the form $f = \theta \cdot \tanh(\theta x/2\lambda)$. The energy of the unit area of this boundary is given by the formula

$$\sigma = 2A\lambda\theta^3/3. \quad (3)$$

The twin structure of smectic crystals must be observed under the conditions of Helfrich instability at strains noticeably larger than the critical value (see the problem in [1, Sect. 45]: the smectic layer of thickness L bounded by solid walls parallel to the smectic layer is extended along the z axis). At very small tensions $\delta L > \delta L_c = 2\pi\lambda$, when δL is about the smectic period, the homogeneous state becomes unstable with respect to the appearance of a periodic structure in the xy plane with wave-number $k_c = \sqrt{\pi/\lambda L}$. At a much larger strain $\delta L \gg \delta L_c$, a twin structure as that schematically shown in the figure should appear.



The parameters of this structure are determined by minimizing the total energy of twin boundaries ($\theta = \varepsilon$, at the vertical boundaries and $\theta = \varepsilon/2$ at the boundaries of the triangular regions). According to geometric consideration, the angle at the vertex of a triangle is equal to ε , the height H of the triangles is related to the structure period d as $\tan(\varepsilon/2) = d/2H$, and the quantity δL is related to the parameter ε as

$$\delta L = (L - H) \left(\frac{1}{\cos \varepsilon} - 1 \right).$$

The energy density of the structure proposed above is given by the expression

$$\frac{1}{Ld} \left\{ 2 \frac{L-H}{\cos \varepsilon} \sigma(\varepsilon) + 4 \frac{H}{\cos(\varepsilon/2)} \sigma(\varepsilon/2) \right\}. \quad (4)$$

In view of the indicated geometric relationships, at a given tension $\delta L/L$ energy (4) is a function of one parameter ε . When $\delta L \ll L$, angle ε is small. In this case, with the use of result (3), the minimum of energy (4) is found to correspond to $\varepsilon = \sqrt{6\delta L/L}$. In this case, $H = 2L/3$ and

$$d = 2\sqrt{\frac{2L\delta L}{3}}.$$

I am grateful to E.I. Kats for stimulating discussions.

1. L.D. Landau and E.M. Lifshitz, Theory of Elasticity, Pergamon, NY (1986)

Translated by R. Tyapaev

JETP Lett. **116**(4), 587 (2007)